



S14 Patients' perceptions of an exercise programme delivered following discharge from hospital after critical illness (the revive trial)

McDowell, K., Bradley, J., McAuley, D., Blackwood, B., & O'Neill, B. (2016). S14 Patients' perceptions of an exercise programme delivered following discharge from hospital after critical illness (the revive trial). In *Unknown Host Publication* (Vol. 71, pp. A10-A10). BMJ Publishing Group. <https://doi.org/10.1136/thoraxjnl-2016-209333.20>

[Link to publication record in Ulster University Research Portal](#)

Published in:
Unknown Host Publication

Publication Status:
Published (in print/issue): 01/12/2016

DOI:
[10.1136/thoraxjnl-2016-209333.20](https://doi.org/10.1136/thoraxjnl-2016-209333.20)

Document Version
Author Accepted version

General rights
Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

recommendations on dose. Recent studies have suggested that some critically ill patients fail to achieve sufficient plasma antibiotic concentrations to treat infection effectively.²

We determined whether critically-ill patients with respiratory infection achieved pharmacokinetic/pharmacodynamic (PK/PD) targets during antibiotic treatment and investigated factors associated with failure to meet these targets.

Methods This was a subgroup, interim analysis of an ongoing study, ABDose. Participants were adults in intensive care receiving piperacillin-tazobactam or co-amoxiclav for respiratory infection. Demographics and measures of organ function were recorded. Antibiotic concentrations were measured, at steady-state, in plasma at 50% and 100% of the dosing interval. Efficacy of beta-lactam antibiotics is dependent upon time above minimum inhibitory concentration (MIC). We chose PK/PD targets of antibiotic concentration >MIC and a more conservative $>4 \times \text{MIC}$ of likely pathogen or microbiological isolate (when available). These targets have been used previously.² During 28-day follow up, need for additional antibiotics was recorded.

Results 24 participants (median age 61, IQR [50–70] years), received co-amoxiclav ($n = 7$), piperacillin-tazobactam ($n = 15$) or both ($n = 2$). At 100% of the dosing interval, 12 achieved plasma antibiotic concentrations >MIC and 8 achieved $>4 \times \text{MIC}$. Participants who did not achieve PK/PD targets were younger (48 [39–59] years vs 68 [61–80] years, $p = 0.002^*$) and had a higher eGFR ($131 \pm 58 \text{ ml/min/1.73m}^2$ vs $64 \pm 28 \text{ ml/min/1.73m}^2$, $p = 0.004^*$) than those who did. Antibiotic concentrations were correlated with age and negatively correlated with eGFR (Figure 1). All participants failing to achieve antibiotic concentrations $>4 \times \text{MIC}$ at 100% of the dosing interval required further courses of antibiotics during follow-up compared to 50% of patients achieving this target ($p = 0.02^*$).

Conclusion In critically-ill patients with respiratory infection, uniform dosing of beta-lactam antibiotics does not consistently achieve PK/PD targets required for optimal efficacy. Younger patients, with better renal function may be under-dosed. These interim findings identify a need for further work to determine whether personalised dynamic dosing regimens could improve outcomes for patients with severe respiratory infection. Population PK modelling and further covariate analysis is planned within ABDose.

REFERENCES

- 1 Mayr FB, et al. Epidemiology of severe sepsis. *Virulence* 2014;**5**:4–11.
- 2 Roberts JA, et al. DALI: defining antibiotic levels in intensive care unit patients: are current β -lactam antibiotic doses sufficient for critically ill patients?. *Clin Infect Dis* 2014;**58**:1072–83.

S14

PATIENTS' PERCEPTIONS OF AN EXERCISE PROGRAMME DELIVERED FOLLOWING DISCHARGE FROM HOSPITAL AFTER CRITICAL ILLNESS (THE REVIVE TRIAL)

¹K McDowell, ²JM Bradley, ³DF McAuley, ²B Blackwood, ¹B O'Neill. ¹Centre for Health and Rehabilitation Technologies, Ulster University, Newtownabbey, UK; ²School of Medicine, Dentistry and Biomedical Sciences, Queens University, Belfast, UK; ³School of Medicine, Dentistry and Biomedical Sciences, Queens University and Regional Intensive Care Unit, Royal Victoria Hospital and Clinical Trials Unit, Belfast, UK

10.1136/thoraxjnl-2016-209333.20

Introduction The REVIVE RCT investigated the effectiveness of an individually tailored (personalised) exercise programme for patients discharged from hospital after critical illness.¹ By

including qualitative methods, we aimed to explore their perceptions of engaging in the 6 week programme to facilitate a better understanding of the intervention and trial outcomes.

Methods Patients allocated to the exercise group were invited to participate in semi-structured interviews following their final outcome assessment (6 months following randomisation). Interviews were conducted by a trained member of the research team not involved in the intervention. Interviews were audio recorded, transcribed verbatim and content analysis used to explore themes arising from the data.

Results Of 30 patients allocated to the exercise group 21 completed interviews. Seven core themes were identified (1) sequelae of critical illness and critical care recovery; (2) satisfaction and endorsement of the exercise programme; (3) beneficial impact of the exercise programme on physical and psychological health; (4) facilitators of beneficial impact; (5) barriers to beneficial impact; (6) challenges to continuing exercise; (7) contrasting views on outcome measures.

Patients provided insight into the physical and mental sequelae they experienced following critical illness. There was a strong sense of patients' need for the exercise programme and its importance for their recovery following discharge home. The programme was described as invaluable, and provided feelings of motivation and hope. Key facilitators of beneficial impact included supervision, tailoring of the exercises to personal needs, and the manual. Barriers to the beneficial impact of the programme included poor mental health, existing physical limitations and lack of motivation. Patients' views of the questionnaires and performance based outcome measures in the REVIVE trial varied. Many patients were unsure about what would be the best way of measuring how the programme affected their health.

Conclusion The benefits of physical rehabilitation programmes, needs to be counterbalanced against patients' mental health status post-ICU and any pre-admission limitations, if they are to be successful. Including this qualitative component improved our understanding of the mechanisms underpinning the impact of the programme and how programmes should be evolved for future trials.

REFERENCE

- 1 O'Neill B, et al. Effectiveness of a programme of exercise on physical function in survivors of critical illness following discharge from the ICU: study protocol for a randomised controlled trial (REVIVE). *Trials* 2014;**15**:146.

S15

CHANGES IN PERIOPERATIVE ARDS WITH TIME: A COMPARISON OF TWO TRIALS

¹PA Howells, ¹KA Aldridge ¹RCA Dancer, ²O Tucker, ¹DR Thickett. ¹Institute of Inflammation and Ageing, University of Birmingham, Birmingham, UK; ²Department of Upper Gastrointestinal Surgery, Heart of England NHS Foundation Trust, Birmingham, UK

10.1136/thoraxjnl-2016-209333.21

Introduction and objective The BALTI-Prevention Trial¹ translational sub-study (recruitment completed in 2011) and VINDALOO Trial² (recruitment completed in 2015) both used oesophagectomy as a model for investigation of the pharmacological prevention of ARDS. The VINDALOO trial showed a lower ARDS incidence independent of the agents evaluated. Our objective was to characterise this difference.

Methods Databases from both trials were available and additional information was obtained retrospectively from hospital records. Analysis was performed using appropriate statistical tests.

THORAX

S14 Patients' perceptions of an exercise programme delivered following discharge from hospital after critical illness (the revive trial)

K McDowell, JM Bradley, DF McAuley, B Blackwood and B O'Neill

Thorax 2016 71: A10

doi: 10.1136/thoraxjnl-2016-209333.20

Updated information and services can be found at:

http://thorax.bmj.com/content/71/Suppl_3/A10.1

Email alerting service

These include:

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Adult intensive care](#) (179)
[Clinical trials \(epidemiology\)](#) (557)
[Sports and exercise medicine](#) (92)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>